

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Haque, Md Mozzammel ORCID logoORCID: <https://orcid.org/0000-0003-1189-4559> and Bayford, Richard ORCID logoORCID: <https://orcid.org/0000-0001-8863-6385> (2019) Protein misfolding thermodynamics. The Journal of Physical Chemistry Letters, 10 (10) . pp. 2506-2507. ISSN 1948-7185 [Article] (doi:10.1021/acs.jpclett.9b00852)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/26662/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

Protein misfolding thermodynamics

Md Mozzammel Haque^{1, 2 *}, Richard Bayford²

¹Middlesex University, The Burroughs, London, NW4 4BT, UK

²ChemiScience, Ashulia, Dhaka-1344, Bangladesh

*mdmozzammelhaque@gmail.com

Abstract: It is known that protein misfolding is governed by the hydrophobic effect of solutes at hydrophobic amino acid side-chains. The hydrophobic force of non-aqueous solutes acts as a driving force for the spatial rearrangement of protein side-chains, whose structural transitions need to be regulated in both time and space. Smaller hydrophobic solutes exert more effect at protein side-chains, which involves the clustering of proteins into misfolded shapes. The consequences of misfolding are either loss of protein function, gain of toxic function or both. This is a physical process, whose result has been directly linked to a large number of human diseases.

Keywords: Hydrophobic effect, Hydrophobic solutes, Misfolding temperature limit,

In cells, proteins function in an aqueous solution. Electrolytes and other small molecules play important roles in the structure formation and function of proteins. A function of proteins in water can broadly be classified as hydrophilic and hydrophobic. Electrolytes and small molecules are inorganic and organic compounds (e.g. most drugs, size 1nm), which support the function of proteins¹.

Electrolytes dissociate completely in aqueous solution are strongly hydrated, which exert hydrophilic interactions (dipole-dipole, ion-dipole, and ion-ion) on hydrophilic sites (Fig. 1a). The hydrophilic force is generally the combination of electrostatic repulsion and attraction between hydrated/non-hydrated ions. The high dielectric constant of water (78.5) weakens the electrostatic interactions between two ions in solution and charged residues in biopolymers and membranes, to the level of thermal energy. At low ionic strengths, electrostatic repulsion (e.g. common ion effect) is a very long range-acting force, which dominates the hydrophobic force, destabilizes hydrophobic interactions² (protein unfolding). In addition, water on

hydrophobic surfaces that exhibit a hydration force (Fig. 1a), which is a strong short-range repulsive force that acts between water and non-polar molecular surfaces³.

Protein folding in cells occurs through a delicate balance of hydration and hydrophilic forces, which are determined by a heterogeneous anisotropic lipid bilayer. For water-soluble proteins, whose folding is majorly driven by the hydrophobic force⁴.

Weak electrolytes (e.g. organic and inorganic) and non-electrolytes (e.g. organometallics, biomolecules etc.) that dissolve in water as ions/molecules have a property of the hydrophobic effect rather than hydrophilic forces. These molecules/ions bind directly with hydrophobic amino acid side chains and loosen protein backbone hydrogen bonds.

The more electronegative elements hold on to their electrons more tightly, which thus are not readily polarized in water. The high charge (oxidation state) to radius ratio of elements (e.g. Al^{3+} , F^-), which determines its specific attraction for hydrophobic sides, is at the origin of its hydrophobic effects on proteins. The hydrophobic force is more active if all bonds in solutes are symmetrical bonds (e.g. CCl_4) that do not change dipole during the interaction.

The hydrophobic force ξ_{HP} of solutes (“wrong” metal/ions, small, large and macromolecules) can be defined by the following equation:

$$\xi_{\text{HP}} = \frac{k_B T}{\sqrt[3]{\zeta \nabla}}$$

Interactions of such solutes at hydrophobic sites as a whole display collective motions and excitations of energy, which are in the order of thermal energy $k_B T$ (where T = absolute temperature and k_B = Boltzmann constant). The actual hydrophobic force achieved by such solutes is the result of the optimization of shapes, which can be determined by using volume⁵ ∇ (∇ = specific volume x molar mass). ζ is a constant, whose value depends on solute shapes.

$$T_{\text{Misfolding}} = \frac{\Delta H}{\kappa_B \tau}$$

In protein folding, $k_B T$ and ΔH are equal (Fig. 1e), which sets a limit on the amount of the temperature by which the protein can be misfolded (Fig.1f). This misfolding temperature is usually much higher than the absolute temperature, which is associated with the gradient of the misfolded energy surface of a peptide that acts as a hydrophobic force within the configuration space. In addition, high ionic strength results in a larger magnitude for ΔH , and an increase in $T_{\text{Misfolding}}$. However, the hydrophobic force that leads to protein shape-changing is entropic at low temperatures and enthalpic at higher temperatures. Therefore, the enthalpic component of the hydrophobic force contributes to protein misfolding.

Conclusions

Although the mechanism of protein misfolding is not fully understood. This paper suggests that the process of protein misfolding, the hydrophobic force of solutes (e.g. “wrong” metals/ions, “wrong” molecules) can be determined, and is higher than the stiffness of protein backbone hydrogen bonds. This force is a result of features that yield a large enthalpy on misfolding. Furthermore, the consequence of this force is the destabilization of the folded states of proteins as a function of temperature. The temperature dependence hydrophobic interactions have been determined as the misfolding temperature, which changes from protein folding to misfolding. These findings could provide a valuable insight into protein structures, which will open new opportunities for the study of the precise misfolding mechanism which is a key feature of many disorders of humans.

References

- (1) Osada, H. *Protein targeting with small molecules: chemical biology techniques and applications*; John Wiley & Sons, Inc.: Hoboken, N.J., 2009.
- (2) Stenner, R. S. *Protein and peptide folding, misfolding and non-folding*; John Wiley & Sons, Inc.: Hoboken, N.J., 2012.
- (3) Disalvo, E. A., *Membrane hydration: the role of water in the structure and function of biological membranes*; Springer: Cham, 2015.
- (4) Petsko, G. A., Ringe, D. *Protein structure and function*; Oxford University Press (1st

Ed): Oxford, 2009.

- (5) Marcus, Y. *Ions in water and biophysical implications: from chaos to cosmos*; Springer: Dordrecht, 2012.
- (6) Scott, R.L., Fernandez, A. *A mathematical approach to protein biophysics*; Springer: Cham, 2017.
- (7) Kuriyan, J., Konforti, B., Wemmer, D. *The molecules of life: physical and chemical principles*; Garland Science: N.Y., 2012.
- (8) Bruns, C. J., Stoddart, J. F. *The nature of the mechanical bond: from molecules to machines*; John Wiley & Sons, Inc.: Hoboken, N.J., 2016.
- (9) Huddler, D., Zartler, E. R. *Applied biophysics for drug discovery*; John Wiley & Sons, Inc.: Chichester, West Sussex, 2017.